

Description

Monitoring Matters: Back-to-Basics: Why Maternal Observations Matter

Samuel Todd – RM, BSc (Hons), PGDip, Consultant Midwife

Social @sjtodd16

Published in The Practising Midwife Volume 29 Issue 5 September 2026, Page 13

https://doi.org/10.55975/MNGP7821

Introduction

Maternal observations throughout pregnancy play a vital role in detecting potential complications early and promoting positive pregnancy outcomes. In essence, they are far more than routine measurements and clinical checkpoints. They provide a vital window into the health and wellbeing of both mother and baby, enabling healthcare professionals to identify emerging concerns, respond proactively and support safer pregnancy outcomes. From the earliest antenatal appointment through to birth, consistent and accurate observation form the foundation of informed clinical decision-making.

Fundamental component

Maternal monitoring is a fundamental component of midwifery care. It encompasses the systematic assessment of physical, psychological and social wellbeing. Asking simple but meaningful questions can provide valuable insight into a person's health, circumstances and support needs.[1]

Effective monitoring includes planned and repeated assessment through clinical observations and screening tests that help identify emerging complications, recognise deterioration and inform timely, proportionate escalation of care.[2]

As maternity services continue to evolve, with increasingly sophisticated technologies and more complex care pathways, it can be easy to lose sight of the fundamentals of midwifery practice. Yet some of the most important opportunities to detect maternal deterioration still arise from two simple assessments performed routinely throughout pregnancy: blood pressure measurement and urinalysis.



Understanding physiological adaptation

Pregnancy is characterised by profound physiological change.[3] Increases in blood volume and cardiac output, alongside reductions in vascular resistance, are essential adaptations that support foetal growth and development. However, these changes can also make it challenging to distinguish normal physiology from the early signs of pathology.

Accurate BP measurement therefore remains a cornerstone of antenatal assessment.[1,2] Importantly, observations should never be considered in isolation. Understanding an individual's baseline and recognising patterns or trends over time is more clinically meaningful than any single reading.[4] Reliable assessment also depends on getting the basics right, including correct cuff selection, appropriate positioning and adherence to best-practice measurement techniques.

The continuing importance of urinalysis

The value of urinalysis should be equally recognised. Its true value lies in the clinical insights it provides. Although often viewed as a routine element of antenatal care, it provides important clinical information that can contribute significantly to decision-making.

Proteinuria, when assessed alongside BP and other symptoms, may indicate the development of pre-eclampsia. Urinalysis may also identify urinary tract infection, glycosuria or ketonuria, prompting further investigation and management where necessary.[1,2] When interpreted alongside maternal observations, symptoms and clinical judgement, urinalysis helps build a picture of maternal wellbeing and supports the early recognition of complications.

Ultimately, technology and innovation will continue to shape maternity care, but the foundations of safe practice remain unchanged. Careful observation, effective communication and the consistent application of fundamental assessments continue to be among the most powerful tools midwives have to keep women safe throughout pregnancy. **TPM**

Five principles for maternal monitoring

1. Know the individual's history, risks and baseline.
2. Use the correct equipment and measurement technique.
3. Interpret observations as trends, not isolated values.
4. Consider symptoms and the whole clinical picture.
5. Document, communicate and escalate concerns promptly.

References

1. WHO recommendations on antenatal care for a positive pregnancy experience. WHO. https://www.who.int/publications/i/item/9789241549912. Published November 28, 2016.
2. Antenatal care. Nice guideline (NG201). NICE. https://www.nice.org.uk/ guidance/ng201. Published August 19, 2021.
3. Murray I, Hassall J. Change and adaptation in pregnancy. In: Marshall JE, Raynor MD, eds. Myles Textbook for Midwives. 17th ed. Elsevier; 2020.
4. Loerup L, Pullon RM, Birks J, et al. Trends of blood pressure and heart rate in normal pregnancies: a systematic review and meta-analysis. BMC Med. 2019;17(1):167. doi:10.1186/s12916-019-1399-1.
5. Knight M, Bunch K, Tuffnell D, Shakespeare J, Patel R, Kurinczuk JJ, eds. Saving Lives, Improving Mothers' Care: Lessons Learned to Inform Maternity Care From the UK and Ireland Confidential Enquiries Into Maternal Deaths and Morbidity 2021–2023. National Perinatal Epidemiology Unit, University of Oxford. https://www.npeu.ox.ac.uk/assets/downloads/mbrrace-uk/ reports/maternal-report-2025/MBRRACE-UK%20Maternal%20Report%20 2025%20-%20Main%20v4.0.pdf. Updated Aug 04, 2026.

Category

1. Editorial
2. Features
3. TPM Articles

4. TPM Journal

Tags

1. 2026
2. fetal monitoring
3. midwifery practice

Date Created
1st September 2026